

**REQUIRED INFORMATION
TO BE READ BY PARENT
AND STUDENT**

**CONCUSSION & CARDIAC
ARREST FORM AT THE
END OF THE DOCUMENT.**

**PRINT, READ, SIGN AND
RETURN THIS FORM (pg.
8) ONLY.**

CONCUSSION FACT SHEET FOR PARENTS



WHAT IS A CONCUSSION?

A concussion is a type of traumatic brain injury. Concussions are caused by a bump or blow to the head. Even a “ding,” “getting your bell rung,” or what seems to be a mild bump or blow to the head can be serious.


You can’t see a concussion. Signs and symptoms of concussion can show up right after the injury or may not appear or be noticed until days or weeks after the injury. If your child reports any symptoms of concussion, or if you notice the symptoms yourself, seek medical attention right away.

WHAT ARE THE SIGNS AND SYMPTOMS OF CONCUSSION?

If your child has experienced a bump or blow to the head during a game or practice, look for any of the following signs of a concussion:

SYMPTOMS REPORTED BY ATHLETE:

- Headache or “pressure” in head
- Nausea or vomiting
- Balance problems or dizziness
- Double or blurry vision
- Sensitivity to light
- Sensitivity to noise
- Feeling sluggish, hazy, foggy, or groggy
- Concentration or memory problems
- Confusion
- Just not “feeling right” or is “feeling down”



SIGNS OBSERVED BY PARENTS/ GUARDIANS:

- Appears dazed or stunned
- Is confused about assignment or position
- Forgets an instruction
- Is unsure of game, score, or opponent
- Moves clumsily
- Answers questions slowly
- Loses consciousness (even briefly)
- Shows mood, behavior, or personality changes

[INSERT YOUR LOGO]



DANGER SIGNS

Be alert for symptoms that worsen over time. Your child or teen should be seen in an emergency department right away if s/he has:

- One pupil (the black part in the middle of the eye) larger than the other
- Drowsiness or cannot be awakened
- A headache that gets worse and does not go away
- Weakness, numbness, or decreased coordination
- Repeated vomiting or nausea
- Slurred speech
- Convulsions or seizures
- Difficulty recognizing people or places
- Increasing confusion, restlessness, or agitation
- Unusual behavior
- Loss of consciousness (even a brief loss of consciousness should be taken seriously)

WHAT SHOULD YOU DO IF YOU THINK YOUR CHILD HAS A CONCUSSION?

1. **SEEK MEDICAL ATTENTION RIGHT AWAY**
A health care professional will be able to decide how serious the concussion is and when it is safe for your child to return to regular activities, including sports.
2. **KEEP YOUR CHILD OUT OF PLAY.**
Concussions take time to heal. Don't let your child return to play the day of the injury and until a health care professional says it's OK. Children who return to play too soon - while the brain is still healing - risk a greater chance of having a second concussion. Repeat or later concussions can be very serious. They can cause permanent brain damage, affecting your child for a lifetime.
3. **TELL YOUR CHILD'S COACH ABOUT ANY PREVIOUS CONCUSSION.**
Coaches should know if your child had a previous concussion. Your child's coach may not know about a concussion your child received in another sport or activity unless you tell the coach.

HOW CAN YOU HELP YOUR CHILD PREVENT A CONCUSSION OR OTHER SERIOUS BRAIN INJURY?

- Ensure that they follow their coach's rules for safety and the rules of the sport.
- Encourage them to practice good sportsmanship at all times.
- Make sure they wear the right protective equipment for their activity. Protective equipment should fit properly and be well maintained.
- Wearing a helmet is a must to reduce the risk of a serious brain injury or skull fracture.
 - However, helmets are not designed to prevent concussions. There is no "concussion-proof" helmet. So, even with a helmet, it is important for kids and teens to avoid hits to the head.

HOW CAN I HELP MY CHILD RETURN TO SCHOOL SAFELY AFTER A CONCUSSION?

Children and teens who return to school after a concussion may need to:

- Take rest breaks as needed
- Spend fewer hours at school
- Be given more time to take tests or complete assignments
- Receive help with schoolwork
- Reduce time spent reading, writing, or on the computer

Talk with your child's teachers, school nurse, coach, speech-language pathologist, or counselor about your child's concussion and symptoms. As your child's symptoms decrease, the extra help or support can be removed gradually.



JOIN THE CONVERSATION  www.facebook.com/CDCHeadsUp

TO LEARN MORE GO TO >> WWW.CDC.GOV/CONCUSSION

Content Source: CDC's Heads Up Program. Created through a grant to the CDC Foundation from the National Operating Committee on Standards for Athletic Equipment (NOCSAE).

A FACT SHEET FOR Athletes



This sheet has information to help you protect yourself from concussion or other serious brain injury and know what to do if a concussion occurs.

WHAT IS A CONCUSSION?

A concussion is a brain injury that affects how your brain works. It can happen when your brain gets bounced around in your skull after a fall or hit to the head.

What Should I Do If I Think I Have a Concussion?

Report It.

Tell your coach and parent if you think you or one of your teammates may have a concussion. You won't play your best if you are not feeling well, and playing with a concussion is dangerous. Encourage your teammates to also report their symptoms.



Get Checked Out by a Doctor.

If you think you have a concussion, do not return to play on the day of the injury. Only a doctor or other healthcare provider can tell whether you have a concussion and when it's OK to return to school and play.



Give Your Brain Time to Heal.

Most athletes with a concussion get better within a couple of weeks. For some, a concussion can make everyday activities, such as going to school, harder. You may need extra help getting back to your normal activities. Be sure to update your parents and doctor about how you are feeling.



GOOD TEAMMATES KNOW:
IT'S BETTER TO MISS ONE GAME THAN THE WHOLE SEASON.



cdc.gov/HEADSUP

How Can I Tell If I Have a Concussion?

You may have a concussion if you have any of these symptoms after a bump, blow, or jolt to the head or body:

-  **Get a headache**
-  **Feel dizzy, sluggish, or foggy**
-  **Are bothered by light or noise**
-  **Have double or blurry vision**
-  **Vomit or feel sick to your stomach**
-  **Have trouble focusing or problems remembering**
-  **Feel more emotional or “down”**
-  **Feel confused**
-  **Have problems with sleep**

A concussion feels different to each person, so it's important to tell your parents and doctor how you feel. You might notice concussion symptoms right away, but sometimes it takes hours or days until you notice that something isn't right.


How Can I Help My Team?

Protect Your Brain.

All your teammates should avoid hits to the head and follow the rules for safe play to lower chances of getting a concussion.

Be a Team Player.

If one of your teammates has a concussion, tell them that they're an important part of the team and they should take the time they need to get better.



The information provided in this document or through linkages to other sites is not a substitute for medical or professional care. Questions about diagnosis and treatment for concussion should be directed to a physician or other healthcare provider.

Revised January 2019

To learn more,
go to cdc.gov/HEADSUP



SUDDEN CARDIAC ARREST

A Fact Sheet for Parents

FACTS

Sudden cardiac arrest is a rare, but tragic event that claims the lives of approximately 500 athletes each year in the United States. Sudden cardiac arrest can affect all levels of athletes, in all sports, and in all age levels. The majority of cardiac arrests are due to congenital (inherited) heart defects. However, sudden cardiac arrest can also occur after a person experiences an illness which has caused an inflammation to the heart or after a direct blow to the chest.

WARNING SIGNS

There may not be any noticeable symptoms before a person experiences loss of consciousness and a full cardiac arrest (no pulse and no breathing).

Warning signs can include a complaint of:

- Chest Discomfort
- Unusual Shortness of Breath
- Racing or Irregular Heartbeat
- Fainting or Passing Out

EMERGENCY SIGNS – Call EMS (911)

If a person experiences any of the following signs, call EMS (911) immediately:

- *If an athlete collapses suddenly during competition*
- *If a blow to the chest from a ball, puck or another player precedes an athlete's complaints of any of the warning signs of sudden cardiac arrest*
- *If an athlete does not look or feel right and you are just not sure*

How can I help my child prevent a sudden cardiac arrest?

Daily physical activity, proper nutrition, and adequate sleep are all important aspects of life-long health. Additionally, parents can assist student athletes prevent a sudden cardiac arrest by:

- Ensuring your child knows about any family history of sudden cardiac arrest (onset of heart disease in a family member before the age of 50 or a sudden, unexplained death at an early age)
- Ensuring your child has a thorough pre-season screening exam prior to participation in an organized athletic activity
- Asking if your school and the site of competition has an automatic defibrillator (AED) that is close by and properly maintained
- Learning CPR yourself
- Ensuring your child is not using any non-prescribed stimulants or performance enhancing drugs
- Being aware that the inappropriate use of prescription medications or energy drinks can increase risk
- Encouraging your child to be honest and report symptoms of chest discomfort, unusual shortness of breath, racing or irregular heartbeat, or feeling faint

What should I do if I think my child has warning signs that may lead to sudden cardiac arrest?

1. *Tell your child's coach about any previous events or family history*
2. *Keep your child out of play*
3. *Seek medical attention right away*

SUDDEN CARDIAC ARREST

A Fact Sheet for Student Athletes

FACTS

Sudden cardiac arrest can occur even in athletes who are in peak shape. Approximately 500 deaths are attributed to sudden cardiac arrest in athletes each year in the United States. Sudden cardiac arrest can affect all levels of athletes, in all sports, and in all age levels. The majority of cardiac arrests are due to congenital (inherited) heart defects. However, sudden cardiac arrest can also occur after a person experiences an illness which has caused an inflammation to the heart or after a direct blow to the chest. Once a cardiac arrest occurs, there is very little time to save the athlete, so identifying those at risk before the arrest occurs is a key factor in prevention.

WARNING SIGNS

There may not be any noticeable symptoms before a person experiences loss of consciousness and a full cardiac arrest (no pulse and no breathing).

Warning signs can include a complaint of:

- Chest Discomfort
- Unusual Shortness of Breath
- Racing or Irregular Heartbeat
- Fainting or Passing Out

EMERGENCY SIGNS – Call EMS (911)

If a person experiences any of the following signs, call EMS (911) immediately:

- *If an athlete collapses suddenly during competition*
- *If a blow to the chest from a ball, puck or another player precedes an athlete's complaints of any of the warning signs of sudden cardiac arrest*
- *If an athlete does not look or feel right and you are just not sure*

How can I help prevent a sudden cardiac arrest?

Daily physical activity, proper nutrition, and adequate sleep are all important aspects of life-long health. Additionally, you can assist by:

- Knowing if you have a family history of sudden cardiac arrest (onset of heart disease in a family member before the age of 50 or a sudden, unexplained death at an early age)
- Telling your health care provider during your pre-season physical about any unusual symptoms of chest discomfort, shortness of breath, racing or irregular heartbeat, or feeling faint, especially if you feel these symptoms with physical activity
- Taking only prescription drugs that are prescribed to you by your health care provider
- Being aware that the inappropriate use of prescription medications or energy drinks can increase your risk
- Being honest and reporting symptoms of chest discomfort, unusual shortness of breath, racing or irregular heartbeat, or feeling faint

What should I do if I think I am developing warning signs that may lead to sudden cardiac arrest?

1. *Tell an adult – your parent or guardian, your coach, your athletic trainer or your school nurse*
2. *Get checked out by your health care provider*
3. *Take care of your heart*
4. *Remember that the most dangerous thing you can do is to do nothing*

CONCUSSION and SUDDEN CARDIAC ARREST
ACKNOWLEDGEMENT AND SIGNATURE FORM FOR PARENTS AND STUDENT ATHLETES

Student Athlete's Name (Please Print): _____

Sport Participating In (Current and Potential): _____

School: _____ Grade: _____

IC 20-34-7 and IC 20-34-8 require schools to distribute information sheets to inform and educate student athletes and their parents on the nature and risk of concussion, head injury and sudden cardiac arrest to student athletes, including the risks of continuing to play after concussion or head injury. These laws require that each year, before beginning practice for an interscholastic sport, a student athlete and the student athlete's parents must be given an information sheet, and both must sign and return a form acknowledging receipt of the information to the student athlete's coach.

IC 20-34-7 states that an interscholastic student athlete, in grades 5-12, who is suspected of sustaining a concussion or head injury in a practice or game, shall be removed from play at the time of injury and may not return to play until the student athlete has received a written clearance from a licensed health care provider trained in the evaluation and management of concussions and head injuries, and at least twenty-four hours have passed since the injury occurred.

IC 20-34-8 states that a student athlete who is suspected of experiencing symptoms of sudden cardiac arrest shall be removed from play and may not return to play until the coach has received verbal permission from a parent or legal guardian for the student athlete to return to play. Within twenty-four hours, this verbal permission must be replaced by a written statement from the parent or guardian.

Parent/Guardian - please read the attached fact sheets regarding concussion and sudden cardiac arrest and ensure that your student athlete has also received and read these fact sheets. After reading these fact sheets, please ensure that you and your student athlete sign this form, and have your student athlete return this form to his/her coach.

As a student athlete, I have received and read both of the fact sheets regarding concussion and sudden cardiac arrest. I understand the nature and risk of concussion and head injury to student athletes, including the risks of continuing to play after concussion or head injury, and the symptoms of sudden cardiac arrest.

(Signature of Student Athlete)

(Date)

I, as the parent or legal guardian of the above named student, have received and read both of the fact sheets regarding concussion and sudden cardiac arrest. I understand the nature and risk of concussion and head injury to student athletes, including the risks of continuing to play after concussion or head injury, and the symptoms of sudden cardiac arrest.

(Signature of Parent or Guardian)

(Date)